



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	Books
Search <u>Protein</u> <input type="checkbox"/> for _____								Go Clear
Limits		Preview/Index		History		Clipboard		Details
Display	<u>default</u> <input type="checkbox"/>	Show: <u>20</u> <input type="checkbox"/>	Send to	<u>File</u> <input type="checkbox"/>	Get Subsequence			

☐ **1: I39311. Kruppel-type zinc...**[gi:2136376]

BLink, Domains, Links

LOCUS I39311 572 aa linear PRI 01-DEC-2000
 DEFINITION Kruppel-type zinc finger protein ZNF74 - human.
 ACCESSION I39311
 VERSION I39311 GI:2136376
 DBSOURCE pir: locus I39311;

summary: #length 572 #molecular-weight 64193 #checksum 1376
 ;
 superfamily: zinc finger protein ZFP-36; LIM metal-binding repeat
 homology
 ;
 PIR dates: 06-Sep-1996 #sequence_revision 06-Sep-1996 #text_change
 01-Dec-2000

KEYWORDS

SOURCE Homo sapiens (human)

ORGANISM Homo sapiens

Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.

REFERENCE 1 (residues 1 to 572)

AUTHORS Aubry,M., Marineau,C., Zhang,F.R., Zahed,L., Figlewicz,D.,
 Delattre,O., Thomas,G., de Jong,P.J., Julien,J.P. and Rouleau,G.A.

TITLE Cloning of six new genes with zinc finger motifs mapping to short
 and long arms of human acrocentric chromosome 22 (p and q11.2)

JOURNAL Genomics 13 (3), 641-648 (1992)

MEDLINE 92347859

PUBMED 1639391

REFERENCE 2 (residues 1 to 572)

AUTHORS Aubry,M., Demczuk,S., Desmaze,C., Aikem,M., Aurias,A., Julien,J.P.
 and Rouleau,G.A.

TITLE Isolation of a zinc finger gene consistently deleted in DiGeorge
 syndrome

JOURNAL Hum. Mol. Genet. 2 (10), 1583-1587 (1993)

MEDLINE 94093543

PUBMED 8268910

FEATURES

source

Location/Qualifiers

1..572

/organism="Homo sapiens"

/db_xref="taxon:9606"

Protein

1..572

/product="Kruppel-type zinc finger protein ZNF74"

ORIGIN




```

1 mlenyqnlla lgpplhkpdv ishlergeep wsmqrevprg pcpewelkav psqqggicke
61 epaqepimer plggaqawgr qagalqrsqa apgrtchgl grpveefplr cplfaqqrvp
121 eggplldtrk nvqategtrk aparlcagen astpsepekf pqvrrqrgag agegefcge
181 cgkafrqsss ltlhrrwhsr ekaykcdecg kaftwstnll ehrrhtgek pffcgecgka
241 fschsslnvh qrihtgerpy kcsacekafs cssllsmhlr vhtgekpyrc gecgkafnqr
301 thltrhhrik tgekpyqcs cgkaftchss ltvhekihsg dkpfkcsdce kafnsrslr
361 lhqrthtgek pfkacdcgkg fschayllvh rrihsgekp kcnecgkafs shaylivhrr
421 ihtgekpfdc sqcwkafsch sslimhqrih tgekpykcse cgrafsqnhc likhqkihs
481 eksfkcekcq emfnwsshlt ehqrlhsegk plaiqfnkhl lstyyvpgsl lgagdaglrd

```

// 541 vdpidaldva kllcvppra grnfslgskp rn

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Entrez PubMed Nucleotide Protein Genome Structure PMC Taxonomy Boo

Search for

Limits Preview/Index History Clipboard Details

Show:

☐ 1: 1K5JE. Chain E, The Crys...[gi:16974926]

[BLink](#), [Domains](#), [Links](#)

LOCUS 1K5J_E 124 aa linear VRT 10-OCT-2001

DEFINITION Chain E, The Crystal Structure Of Nucleoplasmin-Core.

ACCESSION 1K5J_E

VERSION 1K5J_E GI:16974926

DBSOURCE pdb: molecule 1K5J, chain 69, release Oct 10, 2001;
deposition: Oct 10, 2001;
class: Chaperone;
source: Mol_id: 1; Organism_scientific: Xenopus Laevis; Gene:
Nucleoplasmin; Expression_system: Escherichia Coli;
Expression_system_common: Bacteria; Expression_system_strain:
Bl21(De3); Expression_system_vector_type: Plasmid;
Expression_system_plasmid: Prk172;
Exp. method: X-Ray Diffraction.

KEYWORDS .

SOURCE Xenopus laevis (African clawed frog)

ORGANISM Xenopus laevis
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Amphibia; Batrachia; Anura; Mesobatrachia; Pipioidea; Pipidae;
Xenopodinae; Xenopus.

REFERENCE 1 (residues 1 to 124)

AUTHORS Dutta,S., Akey,I.V., Dingwall,C., Hartman,K.L., Laue,T.,
Nolte,R.T., Head,J.F. and Akey,C.W.

TITLE The crystal structure of nucleoplasmin-core: implications for
histone binding and nucleosome assembly

JOURNAL Mol. Cell 8 (4), 841-853 (2001)

MEDLINE 21541925

PUBMED 11684019

REFERENCE 2 (residues 1 to 124)

AUTHORS Dutta,S., Akey,I.V., Dingwall,C., Hartman,K.L., Laue,T.,
Nolte,R.T., Head,J.F. and Akey,C.W.

TITLE Direct Submission

JOURNAL Submitted (10-OCT-2001)

COMMENT Revision History:
NOV 21 1 Typographical
NOV 1 1 Initial Entry.

FEATURES Location/Qualifiers

source 1..124
/organism="Xenopus laevis"
/db_xref="taxon:8355"

SecStr 17..24
/sec_str_type="sheet"
/note="strand 39"

SecStr 28..32
/sec_str_type="sheet"
/note="strand 40"

SecStr 43..49
/sec_str_type="sheet"
/note="strand 41"

SecStr 50..53
/sec_str_type="sheet"
/note="strand 42"
SecStr 60..67
/sec_str_type="sheet"
/note="strand 43"
SecStr 75..82
/sec_str_type="sheet"
/note="strand 44"
SecStr 87..91
/sec_str_type="sheet"
/note="strand 45"
SecStr 94..97
/sec_str_type="sheet"
/note="strand 46"
SecStr 99..106
/sec_str_type="sheet"
/note="strand 47"
SecStr 110..118
/sec_str_type="sheet"
/note="strand 48"


ORIGIN

1 mastvsntsk lēkpvsliwg celneqnktf efkveddeek cehqlalrtv clgdkakdef
61 hivēivtqēē gaēksvpiat lkpsilpmat mvgiēltppv tfrlkagsgp lyisgqhvam
121 ēēdy

//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Dec 22 2003 07:48:22



Real-Time Solutions
Instruments & Reagents

SCIENCE ONLINE SCIENCE MAGAZINE HOME SCIENCE NOW NEXT WAVE STKE/AIDS/SAGE SCIENCE CAREERS E-MARKETPLACE

Institutions: US Patent & Trademark Office | Sign In as Individual | FAQ | Access Rights | Join AAAS

Science magazine

HELP SUBSCRIPTIONS

SIGN IN

AAAS

SEARCH

BROWSE

ORDER THIS ARTICLE

CORRECTIONS AND CLARIFICATIONS

RESEARCH ARTICLES: "Delineation of mRNA export pathways by the use of cell-permeable peptides" by I.-E. Gallouzi and J. A. Steitz (30 Nov. 2001, p. 1895). The sequences of several of the peptides used were reported incorrectly in Fig. 1A. The actual amino acid sequences that were conjugated to AP are as follows, with substitutions indicated in bold, additions denoted by underlining, and positions of amino acids not present in the peptides used indicated by [-]: HNS:

RRFGGPVHHQAQRFRFSPMGVDHMSGLSGVNVP; NES:

[-]QLPPLERLTLD; mNES: [-]QLPPDLRLTLD; and M9:

NNQSSNFGPMKGGNFGGRSSGPYGGGGQYFAKPRNQ[---]. It has been verified that the substitution of L for I [I is present in the HNS sequence of HuR; X. C. Fan, J. A. Steitz, *Proc. Natl. Acad. Sci. U.S.A.* **95**, 15293 (1998)] does not alter the activity of the AP-HNS in the heterokaryon shuttling assay. Similarly, the absence of

NH₂-terminal N and presence of COOH-terminal GGY (as in hnRNP A1) does not alter the activity of AP-M9. The mNES sequence used and reported above is that of the well-characterized NES mutation called M10 [M. H. Malim, S. Bohnlein, J. Hauber, B. R. Cullen, *Cell* **58**, 205 (1989)]; like the misrecorded mNES sequence, it differs from NES in only two amino acids. The scHNS and scM9 sequences originally reported are scrambled versions of the correct HNS and M9 sequences. Nicholas K. Conrad and Angie S. Grech are acknowledged for their work in discovering the errors and repeating the experiments.

► **dEbate:** Submit a response to this article

► Download to Citation Manager

► Alert me when: new articles cite this article

► Search for similar articles in: Science Online

► Search Medline for articles by:

► This article appears in the following Subject Collections: **Corrections**